

3.5.2 Rapid Service/Prediction Centre

Processing Techniques

The algorithm used by the IERS Rapid Service/Prediction Centre for the determination of the quick-look Earth orientation parameters (EOP) is based on a weighted cubic spline with adjustable smoothing fit to contributed observational data (McCarthy & Luzum, 1991a). Contributed data are corrected for possible systematic differences. Biases and rates are determined with respect to the C04 system of the IERS Earth Orientation Centre (EOC). Statistical weighting used in the spline is proportional to the inverse square of the estimated accuracy of the individual techniques. Minimal smoothing is applied, consistent with the estimated accuracy of the observational data.

Weights in the algorithm may be either a priori values estimated by the standard deviation of the residual of the techniques or values based on the internal precision reported by contributors. Estimated accuracies of data contributed to the IERS Rapid Service/Prediction Centre are given in Table 1. These estimates are based on the statistical reports that were generated weekly as a part of the Bulletin A Rapid Service EOP solution for 2003.

Table 1: Estimated accuracies of the techniques in 2003. Units are milliseconds of arc for x , y , $\delta\psi$, $\delta\varepsilon$, dX , and dY and milliseconds of time for UT1-UTC and LOD.

Contributor Information Name, Sample Rate ¹ , Type	Estimated Accuracy					
	x	y	UT1	LOD	$\delta\psi$ (dX)	$\delta\varepsilon$ (dY)
CSR 3-day SLR	0.40	0.43	0.106*			
DUT ² 1 & 3-day SLR	0.34	0.25				
IAA daily SLR	0.14	0.16				
MCC daily SLR	0.16	0.16				
GSFC daily VLBI Int			0.022			
SPBU daily VLBI Int			0.016			
GSFC twice-weekly VLBI	0.07	0.06	0.002		0.4	0.1
IAA ³ twice-weekly VLBI	0.09	0.10	0.004		(0.1)	(0.1)
IVS twice-weekly VLBI	0.04	0.05	0.004		0.4	0.1
USNO twice-weekly VLBI	0.09	0.09	0.006		0.4	0.1
IGS daily Final	0.05	0.03				
IGS daily Rapid	0.03	0.04		0.02*		
USNO daily GPS UT*			0.008*			
EMR daily GPS UT*			0.02*			
USNO daily AAM UT			0.006			

*All satellite techniques provide information on the rate of change of Universal Time contaminated by effects due to unmodeled orbit node motion. VLBI-based results have been used to correct for LOD biases and to minimize drifts in UT estimates.

¹ The sample rate of provided data sets and not the series update rate.

² In July, DUT SLR sample rate changed from 3-days to daily.

³ IAA VLBI celestial pole offsets are in terms of dX/dY using IAU 2000A Nutation Theory.

Operationally, the weighted spline uses as input the epoch of observation, the observed value, and the weight of each individual data point. The software computes the spline coefficients for every data point which are then used to interpolate the Earth orientation parameter time series so that x , y , UT1-UTC, $\delta\psi$, and $\delta\epsilon$ values are computed at the epoch of zero hours UTC for each day. Since the celestial pole offset software is written in terms of $\delta\psi$ and $\delta\epsilon$, the IAA VLBI dX and dY values are converted to $\delta\psi$ and $\delta\epsilon$ for the combination process. The only data points that are excluded from this process are the points whose errors, as reported by the contributors, are greater than three times their average reported precision or those points that have a residual that is more than four times the associated a priori error estimate. Since all of the observations are reported with the effects of sub-daily variations removed, no processing is done to account for these effects (see IERS Gazette No. 13, 30 January 1997).

Table 2. Mean and standard deviation of the differences between the Rapid Service/Prediction Center solutions and IERS Bulletin B and C04 EOP solutions for 2003. Polar motion X and Y values are in milliseconds of arc and UT1-UTC values are in units of milliseconds of time.

	Bulletin A – Bulletin B		Bulletin A – C04	
	Mean	Std. Deviation	Mean	Std. Deviation
Bulletin A Rapid Running Solution (finals.data)				
X	-0.05	0.07	-0.01	0.07
Y	0.04	0.07	0.05	0.07
UT1-UTC	0.001	0.021	-0.004	0.023
Bulletin A Weekly Solution (finals.data)¹				
X	0.00	0.09	-0.02	0.09
Y	-0.04	0.09	-0.05	0.09
UT1-UTC	-0.007	0.064	-0.005	0.066
Bulletin A Daily Solution (finals.daily)				
X	0.00	0.13	-0.02	0.12
Y	-0.06	0.13	-0.08	0.13
UT1-UTC	-0.003	0.075 (0.053) ²	0.001	0.073 (0.050) ²

¹ Statistics computed over the 7-day combination solution period prior to solution epoch.

² Standard deviation computed over entire year, excluding periods with known VLBI intensive data problems.

The uncertainties in the daily values listed in Bulletin A are derived from the quality of the spline fit in the neighborhood of the day in question. Table 2 shows the accuracies of Rapid Service/Prediction Centre's combination solution for the running, the weekly, and the daily products compared to the Bulletin B and C04 series maintained by the IERS EOC at the Paris Observatory. The running solution is the combination solution over the past 365-day period.

3.5.2 Rapid Service/Prediction Centre

The statistics for the running solution at year's end show the agreement between the Bulletin A running combination solution and the Bulletin B/C04 series for the entire year. The comparison of the 53 weekly solutions to the Bulletin B/C04 series presents the statistics of the residuals computed over the new combination results for




Fig. 1: Differences between Daily Bulletin A Rapid Solutions at each daily solution epoch for 2003 and the Earth Orientation Parameters available in Bulletin B and C04 series produced in March of 2004.

the 7-days prior to the solution epoch. Table 2 also presents the statistics for the daily solution differences for the day of the solution epoch. EOP accuracies for the Bulletin A rapid weekly solution for the new combination for the day of the solution run and daily solution at the time of solution epoch are similar and therefore, not shown in the table.

Figure 1 presents the data used in Table 2 for the determination of the Bulletin A daily solution statistics. Figure 1 and Table 2 both indicate that the Centre's UT1-UTC solution accuracies were degraded during early January, mid- to late-July, late October, and December. These degradations were primarily the result of extended periods without VLBI data and highlight the dependence of EOP on VLBI data. The large 0.15 ms shift in the differences in UT1-UTC between mid-July to early August was traced to problems encountered with the incorporation of new global VLBI solutions into the combination. The large residual in PM-x in Figure 1 on 52892 (10 September 2003) occurred as the result of problems encountered in the initial processing of VLBI data. A similar incident appears to be the cause of the spike in PM-y differences on 52905 (23 September 2003). Additional problems with the VLBI intensive experiments occurred between 24 October and 5 November 2003 due to the temporary loss of the Kokee station and the substitution of the Westford station. Overall, the agreement between the Bulletin A solutions and the IERS EOC solutions is quite good.

Prediction Techniques

Polar motion predictions are based on the extrapolation of an annual and semiannual ellipse and a Chandler circle fit to the previous 400 days of observed values of x and y (McCarthy and Luzum, 1991b; Johnson, 2002). The differences between the last observed pole position and rate and those of the curve are computed. These differences are then used to adjust the extrapolated curve by an amount that decreases with the length of the forecast. In February 1998, the near-term polar motion predictions (less than about 30 days) were improved significantly by modifying the transition process from the last observed polar motion result to the long-term predictions. Continuity in the first derivatives was enforced placing great weight on the observed polar motion rate reported by the IGS in their Rapid series. The improvement was most pronounced for the shortest prediction intervals. The procedure for UT1-UTC involves a simple technique of differencing (McCarthy and Luzum, 1991b). All known effects such as leap seconds, solid Earth zonal tides, and seasonal effects are first removed from the observed values of UT1-UTC. Then, to determine a prediction of UT1-UTC n days into the future, $(UT1-TAI)_{n,}$ the smoothed time value from n days in the past, $\langle(UT1R-TAI)_{,-n}\rangle$ is subtracted from the most recent value, $(UT1R-TAI)_0$.

3.5.2 Rapid Service/Prediction Centre

$$(UT1-TAI)_n = 2(UT1R-TAI)_o - \langle (UT1R-TAI) \rangle_n$$

The amount of smoothing used in this procedure depends on the length of the forecast. Short-term predictions with small values of n make use of less smoothing than long-term predictions. Once this value is obtained, it is possible to account for known effects in order to obtain the prediction of UT1-UTC. This process is repeated for each day's prediction.

The very near-term UT1-UTC prediction is strongly influenced by the observed daily Universal Time estimates derived at USNO from the motions of the GPS orbit planes reported by the IGS Rapid service. The IGS estimates for LOD are combined with the GPS-based UT estimates to constrain the UT1 rate of change for the most recent observation.

The near-term to sub-monthly UT1-UTC prediction also makes use of a UT1-like data product derived from the operational NCEP AAM analysis and forecast data (UTAAM). For the 5 days after the latest observation, AAM-based predictions of LOD excitation are combined smoothly with the longer-term UT1 predictions described above.

Errors of the estimates are derived from analyses of the past differences between observations and the published predictions. Formulas published in Bulletin A are used to extend the tabular data. The predictions of $\delta\psi$ and $\delta\epsilon$ are based on the IERS Conventions (McCarthy, 1996; McCarthy and Petit, 2004). Table 3 shows the standard deviation of the differences between the Bulletin A daily solution predictions and the C04 solution for 2003. Initial Centre estimates indicated that the UT1-UTC prediction performance would be improved by 42% at 10 days into the future (Johnson et al., 2003). However, comparisons of the UT1-UTC prediction performance from 2003 to those estimated in 2001 (before UTAAM was introduced) indicated a better than 50% improvement in prediction error at both 10 days and 20 days into the future by the addition of UTAAM to the combination and prediction routine.

Table 3: Standard Deviation of the differences between the Earth Orientation Parameter time series predictions produced by the Bulletin A daily solutions and the C04 solutions for 2003.

Days in Future	PM-X mas	PM-Y mas	UT1-UTC ms
1	.43	.39	.114
5	2.13	1.55	.425
10	4.22	2.29	.882
20	8.34	3.44	2.03
40	14.7	6.53	4.99
90	19.2	16.9	12.5

The predictions of celestial pole offsets (both dX/dY and $\delta\psi/\delta\epsilon$ representations) are based solely on VLBI data. If no new VLBI 24-hour session observations are available, a new rapid combination/prediction of these angles is not determined. Therefore, the predictions of celestial pole offset start before the solution epoch time and the length of the prediction into the future can and does vary in the daily solution files. The differences between the daily Bulletin A predictions and those of the C04 for 2003 are given in Table 4.

Table 4: Standard Deviation of the differences between the Celestial Pole Offset time series predictions produced by the Bulletin A daily solutions and the C04 solutions for 2003.

Days in Future	dX mas	dY mas	$\delta\psi$ mas	$\delta\epsilon$ mas
1	.12	.12	.29	.12
5	.12	.14	.29	.14
10	.14	.15	.35	.16
20	.14	.15	.44	.22
40	.17	.17	.50	.24

Predictions, of UT1-TAI up to 2013 January 1, are given in Table 5 below. They are derived using a prediction algorithm similar to that employed in the Bulletin A predictions of UT1-UTC. Up to twenty years of past observations of UT1-TAI are used. Estimates of the expected one-sigma error for each of the predicted values are also given. These are based on analyses of the past performance of the model with respect to the observations.

Additional information on recent improvements to IERS Bulletin A and the significance for predictions of GPS orbits for real-time users is provided in the paper by Luzum et al. (2001).

Centre Activities in 2003

During 2003 a number of changes occurred that affected the performance of IERS Bulletin A. In January the IAU resolutions with respect to the IAU 2000A Nutation model were introduced into the Centre's products. The `finals2000A.data` and the `finals2000A.daily` solution files were created to comply with these resolutions. These new files contain the nutation in terms of dX/dY with the effects of the free core nutation computed using the IAU2000A Nutation Theory. Due to problems with the software upgrades and the lack of VLBI intensives, the EOP solution for January 2nd was delayed until early January 3rd. In late summer, `finals2000A.all` was added to our list of products. The `finals2000A.all` file is the `finals.all` file converted to the IAU2000A Nutation Theory in the dX/dY paradigm.

3.5.2 Rapid Service/Prediction Centre

Table 5: Predicted values of UT1-TAI, 2004-2013. Note that TT-UT1 can be obtained from this table using the expression $TT-UT1=32.184s-(UT1-TAI)$.

DATE		UT1-TAI (s)	Uncertainty (s)
2004	Jan 1	-32.389	.001
	Apr 1	-32.433	.001
	Jul 1	-32.45	.02
	Oct 1	-32.41	.04
2005	Jan 1	-32.50	.06
	Apr 1	-32.76	.09
	Jul 1	-33.0	.1
	Oct 1	-33.3	.2
2006	Jan 1	-33.5	.2
	Apr 1	-33.8	.3
	Jul 1	-34.0	.4
	Oct 1	-34.2	.4
2007	Jan 1	-34.4	.5
	Apr 1	-34.7	.6
	Jul 1	-34.9	.8
	Oct 1	-35.1	.9
2008	Jan 1	-35.	1.
	Apr 1	-35.	1.
	Jul 1	-35.	1.
	Oct 1	-35	1.
2009	Jan 1	-35.	2.
	Apr 1	-35.	2.
	Jul 1	-35.	2.
	Oct 1	-35.	2.
2010	Jan 1	-36.	2.
	Apr 1	-36.	2.
	Jul 1	-36.	2.
	Oct 1	-36.	2.
2011	Jan 1	-36.	3.
	Apr 1	-36.	3.
	Jul 1	-36.	3.
	Oct 1	-37.	3.
2012	Jan 1	-37.	3.
	Apr 1	-37.	3.
	Jul 1	-37.	4.
	Oct 1	-37.	4.
2013	Jan 1	-37.	4.

On 15 January 2003 two additional VLBI 24-hour solutions were added to the combination solution. These new VLBI solutions are the USNO R1/R4 VLBI solution and the IVS Combination EOPS VLBI solution. The addition of these series to the solution has improved the robustness of the celestial pole offset parameters. These two VLBI series, like the GSFC VLBI series, are checked daily for updates. If an update is available, then the updated series is used in the next daily solution or weekly solution.

In late January the biases and rates that are applied to the VLBI data were adjusted for the new global solution from GSFC, IAA, and USNO. This update resulted in a small correction in the UT1-UTC mean value with respect to the C04 series.

On 11 February 2003 the retrieval and updating of Saint Petersburg University VLBI intensive data was automated and added to the daily solution software package.

On 20 February 2003 the retrieval and updating of IAA VLBI 24-hr session series was automated. With the automation of this VLBI data set, all of the VLBI 24-hr and intensive sessions are currently updated in the daily solution as soon as new updates become available.

On 5 March 2003 the software was modified and code was written to handle the IAA's switch to the dX/dY paradigm to represent the motion of the pole in the celestial reference system. The IAA's dX/dY values are converted to the $\delta\psi/\delta\epsilon$ paradigm for the actual combination process. The routine used for the conversion between dX/dY and $\delta\psi/\delta\epsilon$ is based on the routines available in the IERS Conventions 2003 (McCarthy and Petit, 2004). However, the conventions do not give guidance in the handling of the dX/dY formal errors computed in the files. Therefore, routines were developed to convert these formal errors back to the $\delta\psi/\delta\epsilon$ paradigm.

In June and July new VLBI global solutions required the estimation of new rates and biases for the combination software. The rates and biases for all of the SLR data sets were also re-estimated. These updates resulted in improvements to the PM-x and PM-y time series. Unfortunately, there was a problem with a VLBI intensive UT1-UTC bias correction, which was detected and corrected in early August.

In August the weekly script was modified. New routines were added to automate the retrieval of the SLR data from the mail server. This modification reduced the weekly processing time.

In late October the rates and biases were recomputed for a few data sets.

Currently, the Centre is investigating ways to improve the estimation of biases between the different Analysis Centre datasets and the C04 series.

Availability of Rapid Service

The data available from the IERS Rapid Service/ Prediction Centre consist mainly of the data used in the IERS Bulletin A. These data include: x , y , UT1-UTC, dX and dY from IAA VLBI; x , y , UT1-UTC, $\delta\psi$ and $\delta\epsilon$ from GSFC VLBI; x , y , UT1-UTC, $\delta\psi$ and $\delta\epsilon$ from USNO VLBI; x , y , UT1-UTC, $\delta\psi$ and $\delta\epsilon$ from IVS combination VLBI; UT1-UTC from Saint Petersburg University 1-day Intensives; UT1-UTC from GSFC 1-day Intensives; x , y , UT1-UTC from CSR LAGEOS 3-day SLR; x , y from Delft University of Technology 3-day SLR and later in the year their 1-day solution; x , y from Institute of Applied Astronomy 1-day SLR; x , y from the Russian Mission Control Centre 1-day SLR; x , y , LOD from the International GPS Service; UT from USNO GPS; UT from NRC Canada (EMR) GPS; UT from NCEP AAM; x , y , UT1-UTC, $\delta\psi$, $\delta\epsilon$ from the IERS Rapid Service/Prediction Centre; x , y , UT1-UTC, $\delta\psi$, $\delta\epsilon$ from the IERS Earth Orientation Centre; and predictions of x , y , UT1-UTC from the IERS Rapid Service/Prediction Centre.

In addition to this published information, other data sets are available. These include: UT0-UTC from University of Texas at Austin LLR, UT0-UTC from JPL LLR; UT0-UTC from CERGA LLR; UT0-UTC from JPL VLBI; latitude and UT0-UTC from Washington PZTs 1,3,7; latitude and UT0-UTC from Richmond PZTs 2,6; x and y from CSR LAGEOS 5-day SLR; x and y from Delft 5-day SLR; and x , y , UT1-UTC, $\delta\psi$, and $\delta\epsilon$ from IRIS VLBI.

The data described above are available from the Centre in a number of forms. You may request a weekly machine-readable version of the IERS Bulletin A containing the current ninety day's worth of predictions via electronic mail from

<ser7@maia.usno.navy.mil> or <<http://maia.usno.navy.mil/>>.

Internet users can also direct an anonymous FTP to

<maia.usno.navy.mil>

and change to the ser7 directory where the IERS Bulletin A and more complete databases can be accessed including the daily Bulletin solutions.

World Wide Web access is available at

<<http://maia.usno.navy.mil/>>.

Centre Staff

For most of 2003, the Rapid Service/Prediction staff consisted of the following members:

William Wooden director

Thomas Johnson program manager, research, and software maintenance

Arvid Myers assists in operations, software maintenance, and support

Merri Sue Carter assists in daily operations and support

William Wooden joined the program in late February 2003.

References

- Johnson, T.J., 2002, Rapid Service/Prediction Centre, *IERS Annual Report 2001*, 47-55.
- Johnson, T.J., Luzum, B.J., and Ray, J.R., 2003, Improved near-term UT1R predictions using forecasts of atmospheric angular momentum, *J. Geodynamics*, in review.
- Luzum, B.J., Ray, J.R., Carter, M.S., and Josties, F.J., 2001, Recent Improvements to IERS Bulletin A Combination and Prediction, *GPS Solutions*, **4**(3), 34.
- McCarthy, D.D. and Luzum, B.J., 1991a, Combination of Precise Observations of the Orientation of the Earth, *Bulletin Geodesique*, **65**, 22.
- McCarthy, D.D. and Luzum, B.J., 1991b, Prediction of Earth Orientation, *Bulletin Geodesique*, **65**, 18.
- McCarthy, D.D., 1996, IERS Conventions (1996), *IERS Technical Note No. 21*, Paris Observatory, France.
- McCarthy, D. and G. Petit, 2004, IERS Conventions (2003), *IERS Technical Note No. 32*, Verlag des Bundesamts für Kartographie und Geodäsie, Frankfurt am Main, Germany.

*Thomas Johnson, Merri Sue Carter, Arvid Myers,
William Wooden*